

## Wednesday, November 12

Program of the day:

- Cutting planes — a method to obtain tighter bounds and faster convergence to integer solutions (Wolsey chap. 8)
- Application: branch-and-cut algorithms

1

## Introduction

- branch-and-bound: divide and conquer.
- cutting plane: add inequalities which separate fractional solution from solution space.

Development

- 50's cutting plane (Gomory: simplex, no  $\mathcal{N}(\mathcal{P}$ -hardness)
- 70's tighten formulation in preprocessing
- 80-90's branch-and-cut (Padberg, Rinaldi)

Preprocessing → part of solution process


Definitions

- cuts: valid inequalities
- facets: inequalities defining convex hull

Cuts and facets are redundant for IP formulation  
Tighten formulation for LP relaxation

2

## Cuts and facets



3

## Examples from last lesson

Preprocessing, integer variables

$$\begin{aligned} & \text{maximize} && \dots \\ & \text{subject to} && 7x_1 + 3x_2 - 4x_3 - 2x_4 \leq 1 \\ & && -2x_1 + 7x_2 + 3x_3 + 4x_4 \leq 6 \\ & && \quad - 2x_2 - 3x_3 - 6x_4 \leq -5 \\ & && 3x_1 \quad \quad - 2x_3 \quad \quad \geq -1 \\ & && x \in \mathbb{B}^4 \end{aligned}$$

Generating logical inequalities

From constraint 1 we see that


- if  $x_1 = 1$  and  $x_2 = 1$  then infeasible, thus

$$x_1 + x_2 \leq 1$$

4

### Examples from last lesson

$$\begin{aligned} &\text{maximize } x_1 + x_2 \\ &\text{subject to } -2x_1 + 2x_2 \geq 1 \\ &\quad -8x_1 + 10x_2 \leq 13 \\ &\quad x_1, x_2 \geq 0, \text{ integer} \end{aligned}$$



### Tightening formulation

$$\begin{aligned} -2x_1 + 2x_2 &\geq 1 & -8x_1 + 10x_2 &\leq 13 \\ -x_1 + x_2 &\geq 1/2 & -4x_1 + 5x_2 &\leq 13/2 \\ -x_1 + x_2 &\geq 1 & -4x_1 + 5x_2 &\leq 6 \end{aligned}$$

5

### Motivation

Integer programming problem (IP)

$$\max\{cx : x \in X\}$$

where  $X = \{x : Ax \leq b, x \in \mathbb{Z}_+^n\}$ . Reformulate to

$$\max\{cx : x \in \text{conv}(X)\}$$

For any  $c$ , an optimal solution to LP is also optimal to IP

### Valid inequalities (def. 8.1)

Consider the problem:

$$\begin{aligned} &\text{maximize } f(x) \\ &\text{subject to } x \in X \end{aligned}$$

An inequality

$$\pi x \leq \pi_0$$

is a *valid inequality* for  $X \subseteq \mathbb{R}^n$  if

$$\pi x \leq \pi_0 \text{ for all } x \in X$$

6

### Characterisation of valid inequalities (sec. 8.3.2)

Consider the problem:

$$\begin{aligned} &\text{maximize } f(x) \\ &\text{subject to } x \in X \end{aligned}$$

where

$$X = \{y \in \mathbb{Z} : y \leq b\}$$

then the inequality

$$y \leq [b]$$

is valid for  $X$

- Simple observation
- Complete characterisation

7

### Overview of cuts

- Chvatal cuts
- Gomory cuts (Modular cuts)
- Chvatal-Gomory cuts
- Disjunctive cuts
- Cover inequalities
- Clique inequalities
- Problem specific cuts

Notice

- Cuts and facets are independent of objective function
- A tight formulation can be used for any objective

8

### Example of Facets

The problem

$$\begin{aligned} &\text{minimize} && 2x_1 + 7x_2 + 2x_3 \\ &\text{subject to} && x_1 + 4x_2 + x_3 \geq 10 \\ &&& 4x_1 + 2x_2 + 2x_3 \geq 13 \\ &&& x_1 + x_2 - x_3 \geq 0 \\ &&& x_1, x_2, x_3 \geq 0, \text{ integer} \end{aligned}$$

has the facets

$$\begin{aligned} x_1 + 4x_2 + x_3 &\geq 10 \\ 2x_1 + x_2 + x_3 &\geq 7 \\ x_1 + x_2 - x_3 &\geq 0 \\ x_1 + 3x_2 + x_3 &\geq 9 \\ 2x_1 + 4x_2 + x_3 &\geq 13 \\ x_1 + x_2 + x_3 &\geq 5 \\ x_1 + 2x_2 &\geq 5 \\ 2x_1 + x_2 &\geq 4 \\ x_1 &\geq 0, \text{ integer} \\ x_2 &\geq 0, \text{ integer} \\ x_3 &\geq 0, \text{ integer} \end{aligned}$$

Using the new formulation we obtain an integer optimal solution by solving the LP-relaxed problem. (For any objective function).

9

### Chvátal Cuts

Valid inequalities for a pure IP-model (minimization)

- 1 Add constraints, using suitable multipliers
- 2 Divide through by a common coefficient factor
- 3 Round up right-hand-side to the next integer

### Example

$$\begin{aligned} &\text{minimize} && 2x_1 + 7x_2 + 2x_3 \\ &\text{subject to} && x_1 + 4x_2 + x_3 \geq 10 && (1) \\ &&& 4x_1 + 2x_2 + 2x_3 \geq 13 && (2) \\ &&& x_1 + x_2 - x_3 \geq 0 && (3) \\ &&& x_1 \geq 0 && (4) \\ &&& x_2 \geq 0 && (5) \\ &&& x_3 \geq 0 && (6) \\ &&& x_1, x_2, x_3 \text{ integer} \end{aligned}$$

1 times (2) is

$$4x_1 + 2x_2 + 2x_3 \geq 13$$

divide by two

$$2x_1 + x_2 + x_3 \geq 6\frac{1}{2}$$

left hand side is integral, thus round up right-hand

$$2x_1 + x_2 + x_3 \geq 7$$

10

### Example (continued)

$$\begin{aligned} &\text{minimize} && 2x_1 + 7x_2 + 2x_3 \\ &\text{subject to} && x_1 + 4x_2 + x_3 \geq 10 && (1) \\ &&& 4x_1 + 2x_2 + 2x_3 \geq 13 && (2) \\ &&& x_1 + x_2 - x_3 \geq 0 && (3) \\ &&& x_1 \geq 0 && (4) \\ &&& x_2 \geq 0 && (5) \\ &&& x_3 \geq 0 && (6) \\ &&& x_1, x_2, x_3 \text{ integer} \end{aligned}$$

Facets

$$\begin{aligned} x_1 + 4x_2 + x_3 &\geq 10 && (a) \\ 2x_1 + x_2 + x_3 &\geq 7 && (b) \\ x_1 + x_2 - x_3 &\geq 0 && (c) \\ x_1 + 3x_2 + x_3 &\geq 9 && (d) \\ 2x_1 + 4x_2 + x_3 &\geq 13 && (e) \\ x_1 + x_2 + x_3 &\geq 5 && (f) \\ x_1 + 2x_2 &\geq 5 && (g) \\ 2x_1 + x_2 &\geq 4 && (h) \\ x_1, x_2, x_3 &\geq 0, \text{ integer} \end{aligned}$$

Obtained as

- (d) :  $5 \times (1), 1 \times (b), 1 \times (6)$ , divide 7
- (f) :  $1 \times (1), 3 \times (b), 3 \times (6)$ , divide 7
- (g) :  $4 \times (1), 1 \times (b), 5 \times (3)$ , divide 11
- (h) :  $1 \times (b), 1 \times (6), 1 \times (4)$ , divide 2
- (e) :  $3 \times (d), 1 \times (b), 3 \times (g)$ , divide 4

11

### Chvatal-Gomory cuts (p. 119)

$$\begin{aligned} &\text{maximize} && \sum_{j=1}^n c_j x_j \\ &\text{subject to} && \sum_{j=1}^n a_{1j} x_j \leq b_1 \\ &&& \vdots \\ &&& \sum_{j=1}^n a_{mj} x_j \leq b_m \\ &&& x_j \in \mathbb{Z}_+, \quad j = 1, \dots, n \end{aligned}$$

1 Take a linear combination of the constraints

$$\sum_{j=1}^n \left( \sum_{i=1}^m u_i a_{ij} \right) x_j \leq \left( \sum_{i=1}^m u_i b_i \right)$$

in short

$$\sum_{j=1}^n a'_j x_j \leq b'$$

2 Since  $x \geq 0$  implies  $\sum_{j=1}^n (a'_j - \lfloor a'_j \rfloor) x_j \geq 0$  we have

$$\sum_{j=1}^n \lfloor a'_j \rfloor x_j \leq b'$$

3 Since  $x_j \in \mathbb{Z}_+$  implies  $\lfloor a'_j \rfloor x_j \in \mathbb{Z}$  we get

$$\sum_{j=1}^n \lfloor a'_j \rfloor x_j \leq \lfloor b' \rfloor$$

12

**Chvatal-Gomory (Theorem 8.4)**

$X = \{x : Ax \leq b, x \in \mathbb{Z}_+^n\}$

Every valid inequality for  $X$  can be obtained by applying the Chvatal-Gomory procedure a finite number of times.

**Notice**


- No stronger inequalities than Chvatal-Gomory exists.
- Even the facet constraints can be generated as Chvatal-Gomory cuts.
- No constructive (polynomial) algorithm for how the linear combination of constraints should be chosen.
- In practice, the derivation of Chvatal-Gomory cuts must rely on specific features of a given application.

Gomory cuts is a systematical way of deriving cutting planes.

**Only 0-1 case**

All bounded integer variables can be expressed as sum of binary variables.

The set  $X = P \cap \mathbb{Z}^n$



$$P = \{x \in \mathbb{R}^n : Ax \leq b, 0 \leq x \leq 1\} \neq \emptyset$$

$$X = P \cap \mathbb{Z}^n$$

Nemhauser and Wolsey, Proposition 1.1 page 208:

If inequality  $\pi x \leq \pi_0$  is valid for  $P$  then it can be obtained as a C-G cut

(\*)

**Proof (0-1 case)**

Assume that

$$\pi x \leq \pi_0 \text{ where } \pi, \pi_0 \text{ integers}$$

is a valid inequality for  $X$ . We will show that this inequality can be obtained by using the C-G procedure a finite number of times.

- Step 1: Find a large number  $t \in \mathbb{Z}_+$  such that

$$\pi x \leq \pi_0 + t$$

is a valid C-G inequality

- Step 2: Prove that if

$$\pi x \leq \pi_0 + \tau + 1$$

for  $\tau \in \mathbb{Z}_+$  is a C-G inequality for  $X$  then also

$$\pi x \leq \pi_0 + \tau$$

is a C-G inequality for  $X$ .

- Step 3: Use step 2 for  $\tau = t - 1, \dots, 0$  each time getting a new C-G inequality

(Proof by induction)

**Step 1**

The inequality

$$\pi x \leq \pi_0 + t$$

is valid for  $P$  for some  $t \in \mathbb{Z}_+$ .

**Proof**

We have the inequality

$$x \leq 1$$

derive C-G inequality using multipliers  $u = \pi$

$$\pi x \leq \pi 1$$

choosing  $t = \pi 1 - \pi_0$  ( $\pi, \pi_0$  is integer) we get the form

$$\pi x \leq \pi 1 = \pi_0 + t$$

for some  $t \in \mathbb{Z}_+$

## Step 2

Difficult part (not proved completely)

- a) Prove that if  $\pi x \leq \pi_0 + \tau + 1$  with  $\tau \in \mathbb{Z}_+$  is a C-G cut then

$$\pi x \leq \pi_0 + \tau + \sum_{j \in N^0} x_j + \sum_{j \in N^1} (1 - x_j)$$

is a C-G inequality for  $X$  for every partition  $(N^0, N^1)$  of  $N = \{1, \dots, n\}$ .

- b) Use partitionings  $(T^0 \cup \{n\}, T^1)$  and  $(T^0, T^1 \cup \{n\})$  to obtain a new inequality for  $(T^0, T^1)$ .
- c) Derive all valid inequalities for partitionings of  $N' = \{1, \dots, n-1\}$
- d) Repeating this procedure  $n$  times implies that we eliminate the sums on the right side and thus

$$\pi x \leq \pi_0 + \tau$$

is a C-G cut

Time complexity

- part (c) takes  $O(2^n)$ ,  
part (d) is performed  $n$  times,  
in total  $O(n2^n)$
- we run Step 2  $O(t)$  times, thus in total  $O(tn2^n)$ .

17

## Step 2, a)

An inequality is valid for  $P$  if it is valid for all vertices  $\{x^1, \dots, x^m\}$  of  $P$

Assume that  $\pi x \leq \pi_0 + \tau + 1$  with  $\tau \in \mathbb{Z}_+$  is a valid cut. Let  $(N^0, N^1)$  be any partitioning of  $N = \{1, \dots, n\}$ . Consider a vertex  $x^k$  of  $P$

- $x^k$  integer: then  $\pi x^k \leq \pi_0$  (since  $\pi x \leq \pi_0$  valid for  $X$ )
- $x^k$  fractional: exists  $\epsilon > 0$  such that

$$\epsilon^k \leq \sum_{j \in N^0} x_j^k + \sum_{j \in N^1} (1 - x_j^k)$$

Choose  $\alpha = \min_{x^k \text{ vertex in } P} \epsilon^k$

Using  $M \geq (\tau + 1)/\alpha$ , we have

$$\tau + 1 \leq M\alpha \leq M \left( \sum_{j \in N^0} x_j + \sum_{j \in N^1} (1 - x_j) \right)$$

adding  $\pi_0$  at both sides we get valid inequality for  $P$

$$\pi x \leq \pi_0 + \tau + 1 \leq \pi_0 + M \left( \sum_{j \in N^0} x_j + \sum_{j \in N^1} (1 - x_j) \right)$$

Due to (\*) the inequality is a C-G cut.

18

## Step 2, a)

We have just shown that the following is a C-G inequality

$$\pi x \leq \pi_0 + M \left( \sum_{j \in N^0} x_j + \sum_{j \in N^1} (1 - x_j) \right)$$

By assumption we had the C-G inequality

$$\pi x \leq \pi_0 + \tau + 1$$

use weights  $1/M$  and  $(M-1)/M$  for the two inequalities getting C-G inequality

$$\pi x \leq \pi_0 + \tau + \sum_{j \in N^0} x_j + \sum_{j \in N^1} (1 - x_j)$$

19

## Step 2, b)

Use partitions  $(T^0 \cup \{n\}, T^1)$  and  $(T^0, T^1 \cup \{n\})$

$$\pi x \leq \pi_0 + \tau + \sum_{j \in T^0 \cup \{n\}} x_j + \sum_{j \in T^1} (1 - x_j)$$

and

$$\pi x \leq \pi_0 + \tau + \sum_{j \in T^0} x_j + \sum_{j \in T^1 \cup \{n\}} (1 - x_j)$$

using multipliers  $1/2$  and  $1/2$  we get C-G inequality

$$\pi x \leq \pi_0 + \tau + \sum_{j \in T^0} x_j + \sum_{j \in T^1} (1 - x_j)$$

20